arXiv:hep-th/9205019v2 26 Jun 1992

IP/BBSR/92-27.

Charged Topological Solitons in 4+1 Dimensional YMCS Theory

C.S.Aulakh
Institute of Physics, Sachivalya Marg
Bhubaneshwar,751005,India

ABSTRACT

We show that when a Chern-Simons term is added to the action of $SU(N)$ $(N \geq 3)$ Yang-Mills theory in 5 dimensions the usual self-dual topological solitons present in the theory necessarily pick up a (topological) electric charge .

Email:aulakh%iopb@shakti.ernet.in

March 1992

IP/BBSR/92-27.

In a recent paper [1] the present author suggested that an examination of the properties of a certain class of co-winding solitons ("syncyclons") generically present in higher dimensional field theories (i.e defined on space-time extended beyond 4 dimensions by a compact space) is called for. Such solitons would appear as point, string or sheet vacuum defects to our three dimensional low energy eyes and might thus signal the presence of otherwise unobservable extra dimensions.

The simplest paradigmatic example of such solitons is furnished by Yang-Mills (YM) theory on $M^4 \times S^1$. The basic solution is nothing but the periodic instanton or *caloron*[3] reinterpreted as a soliton on $M^4 \times S^1$. Classically, such solitons have no (non-abelian) electric charge and, in the Kaluza-Klein (KK) limit $r >> R$ (where R is the radius of S^1 and r the 3-dimensional distance), their fields are those of a non-abelian magnetic dipole together with a scalar potential coming from the extra compoment of the vector potential[1,4]. However, in order to sketch a believable picture of their properties it is necessary to take into account the effects of coupling to fermions and the quantization of global gauge collective coordinates since these can radically change the quantum numbers labelling the soliton states due to fermion number fractionalization[5] and zero point (quantum) rotation in the internal non-abelian space. In particular using standard results on fermion number fractionalization in odd dimensions[5] one finds that the soliton picks up an abelian charge $-\frac{\nu q}{4}$ (ν the Pontryagin index) from each fermion of abelian charge q (Baryon/Lepton number, Electric charge. etc). In addition the expectation value for the charges corresponding to the $U(1)$s in the decomposition $SU(N) \supset SU(2) \times U(1)^{N-2}$ is also proportional to $\nu/4$ (with the soliton trivially embedded in the SU(2) factor). Thus if the gauge group is SU(3) then $< Q^8 >= \frac{-N_f \nu}{4\sqrt{3}}$ where N_f is the number of fermion flavours. The term in the gauge effective action which represents this effect is nothing but the 5 dimensional Chern Simons term times N_f. This term is the low-momentum approximation of the fermion determinant in the gauge field

Email:aulakh%iopb@shakti.ernet.in
March 1992

background. In order to determine the gauge quantum numbers of the soliton one must quantize the collective coordinates of a static solution to the field equations of the YM-Chern Simons (YMCS) system in 5 dimensions. This problem bears a close analogy to that of determining the quantum numbers of the Skyrmeon[7]. The roles of flavor and color are interchanged and the Chern-Simons term is the analog of the WZW term which also arises by integrating out fermions coupled to the Skyrme field . The significant generalization is that the present system is gauge invariant while the Skyrme-WZW Lagrangian has only a global (flavour) invariance. Since the induced charge described above is coupled to a gauge field it follows that here the static solution one must quantize around is a *charged* soliton. In the Skyrme case the effect of the WZW term in the field equations is to produce a deformation of the original soliton without obviously changing its external interactions (as opposed to its flavour and spin quantum numbers). On the contrary, in the present case, the as we shall show, the deformed solution carries a (topological) electric charge. Charged solitons arise in YMCS-Higgs systems in 2+1 dimensions as well[8] where the presence of the CS term allows charged vortex solutions to exist. In 2+1 dimensions the CS term makes the gauge particles massive so that the induced charge couples to a short range force carrier. The integral of the Higgs field charge density is proportional to the quantized flux (divided by 4) so that there too the electric charge is fractional and topological in nature. The existence of charged vortex solutions in 2+1 dimensions can be seen as the analog of the "Witten effect" in 4 dimensions whereby the addition of a CP violating θ term (i.e a Pontryagin density) to the YM-Higgs lagrangian results in charged monopole solutions[6]. One therefore expects that such a conversion of "magnetic" charge neutral solitons to charged ones occurs in all odd dimensions provided the theory without the CS term has a topological soliton solution. Note however that in the present case the magnetic fields are asymptotically *dipolar.*

As a preliminary to the quantization[9] of the collective coordinates of solitons in YM theory on $M^4 \times S^1$, and also because of their intrinsic interest we have studied static solutions of the YMCS system in M^5. The generalization of our results to $M^4 \times S^1$ and the situation in higher odd dimensions will be reported separately[10]. We will show that the above heuristic argument for the existence of charged topological solitons on 5 dimensions is borne out by a detailed analysis of the SU(N), YMCS action:

$$\begin{aligned} S &= \int d^5x \left(\frac{1}{2g_5^2} tr {F_{MN}}^2 + \frac{iN_f}{48\pi^2}\omega_5\right) \\ \omega_5 &= \epsilon_{MNLPQ} \quad tr\left(\partial_M A_N \left(\partial_L A_P A_Q + \frac{3}{2} A_L A_P A_Q\right) + \frac{3}{5} A_M A_N A_L A_P A_Q\right) \end{aligned} \tag{1}$$

The field equations are:

$$D_M F^A_{MN} = -\frac{N_f g_5^2}{128\pi^2} \epsilon_{NMLPQ} \quad tr(\lambda^A F_{ML} F_{PQ}) \tag{2}$$

Our normalizations and definitions are

$$\begin{aligned} A_M &= A^A_M \frac{\lambda^A}{2i} \qquad\qquad tr\lambda^A\lambda^B = 2\delta^{AB} \\ F_{MN} &= F^A_{MN}\frac{\lambda^A}{2i} = \partial_{[M}A_{N]} + [A_M, A_N] \\ A, B, ... &= 1, ..., N^2 - 1 \\ M, N, ... &= 0, 1, 2, 3, 4 \\ \mu, \nu, ... &= 1, 2, 3, 4 \\ a, b, ... &= 1, 2, 3 \\ \hat{a}, \hat{b}, ... &= 4, ..., N^2 - 1 \end{aligned} \tag{3}$$

Here $\{\lambda^A\}$ are the Gell-Mann matrices for SU(N). Our metric convention is $(-++++)$. In the absence of the CS term ($N_f = 0$) we have the usual self dual solutions[11] in the SU(2) Euclidean sector:

Email:aulakh%iopb@shakti.ernet.in

$$
\begin{aligned}
A_0 &= \partial_0 = 0 \\
A_\mu &= -\bar{\eta}_{a\mu\nu}\frac{\tau^a}{2i}\partial_\nu ln\Pi = i\overline{\Sigma}_{\mu\nu}\partial_\nu ln\Pi \\
\Pi^{-1}\partial^2\Pi = 0 \qquad & \qquad \Pi = 1 + \sum_{i=1}^{K}\frac{\rho_i^2}{(x-x_i)^2} \\
\nu &= \frac{1}{32\pi^2}\int d^4x F^a_{\mu\nu}\tilde{F}^a_{\mu\nu} = K
\end{aligned} \tag{4}
$$

where ρ_i, x_i are the scale factors and positions of the K solitons and $\nu = K$ the total winding number. In the above $\{\bar{\eta}_{a\mu\nu}\}$,τ_a, Π are the anti-self-dual 't Hooft symbols, Pauli matrices and "super potential" respectively. Fields outside the $SU(2)$ subalgebra generated by $\{\lambda^a\}$ play no role. The asymptotics are those of a non-abelian magnetic dipole with zero electric charge (since $F_{\mu 0} = 0$). The field energy is

$$
\mathcal{E} = \frac{1}{g_5^2}\int d^4x(\frac{1}{4}{F^a_{\mu\nu}}^2 + \frac{1}{2}{F^a_{\mu 0}}^2) = \frac{8\pi^2\nu}{g_5^2} \tag{5}
$$

We now show that these solutions become charged when the effect of the C.S. term is included. For simplicity let us work with $\nu = K = 1$ and $N = 3$. Generalizations are obvious and will be indicated below. The field equations are

$$
D_M F^A_{M\nu} = \frac{N_f g_5^2}{16\pi^2} \quad tr(\lambda^A F_{\mu 0}\tilde{F}_{\mu\nu}) \tag{6a}
$$

$$
D_\mu F^A_{\mu 0} = -\frac{N_f g_5^2}{64\pi^2} tr(\lambda^A F_{\mu\nu}\tilde{F}_{\mu\nu}) \tag{6b}
$$

It is easy to check that for $A = 8$ the ansatz of (4) gives zero for the l.h.s of eqn.(6b) while the rhs is proportional to the Pontryagin density and is hence nonzero. Thus (4) is no longer an adequate ansatz when $N_f \neq 0$. The required modification is obviously:

Email:aulakh%iopb@shakti.ernet.in

$$\begin{aligned} &A^A_\mu = 0 \qquad\qquad A = 4,5,6,7,8 \\ &A^a_\mu = -\bar{\eta}_{a\mu\nu}\partial_\nu ln\Pi \qquad\qquad \Pi = \Pi(x^2) \\ &A^A_0 = 0 = \partial_0 \qquad A \neq 8 \end{aligned} \tag{7}$$

Spherical symmetry ensures that the r.h.s of the equation for $D_\mu F^a_{\mu 0}$ vanishes as it must for consistency. The only nontrivial equations are then :

$$\begin{aligned} D_\mu F^a_{\mu\nu} &= -\alpha\partial_\mu A^8_0 \tilde{F}^a_{\mu\nu} \\ \partial^2 A^8_0 &= \frac{\alpha}{4} F^a_{\mu\nu}\tilde{F}^a_{\mu\nu} = \frac{\alpha}{2}\partial_\mu\tilde{\omega}_\mu \\ \alpha &= \frac{N_f g_5^2}{32\pi^2\sqrt{3}} \\ \tilde{\omega}_\mu &= \epsilon_{\mu\nu\lambda\gamma}(A^a_\nu\partial_\lambda A^a_\gamma + \frac{\epsilon_{abc}}{3}A^a_\nu A^b_\lambda A^c_\gamma) \end{aligned} \tag{8}$$

Thus the Pontryagin density serves as the charge density in a Poisson equation for the electrostatic potential in the 8 direction. The solution for A^8_0 is simply

$$A^8_0(x) = -\frac{\alpha}{16\pi^2}\int d^4y \frac{1}{(x-y)^2} F^a_{\mu\nu}(y)\tilde{F}^a_{\mu\nu}(y) \tag{9}$$

and also

$$\partial_\mu A^8_0 = \frac{\alpha}{2}(\tilde{\omega}_\mu + \frac{D\hat{x}_\mu}{x^3}) \tag{10}$$

The arbitrarinesss represented by D is crucial to the existence of a charged solution. It is easy to check (assuming that the deformed superpotential has the same leading behaviours as $x \longrightarrow 0$ and $x \longrightarrow \infty$ as before) that the leading behaviour of $\tilde{\omega}_\mu$ as $x \longrightarrow \infty$ is $O(x^{-7})$. Thus to obtain the electric field corresponding to the charge $Q^8 = -\frac{N_f}{4\sqrt{3}}$ we must choose $D = 8$.

The other equation is now

$$D_\mu F^a_{\mu\nu} = -\frac{\alpha^2}{2}(\tilde{\omega}_\mu + \frac{D\hat{x}_\mu}{x^3})\tilde{F}^a_{\mu\nu} \tag{11}$$

Email:aulakh%iopb@shakti.ernet.in March 1992

Due to spherical symmetry the above equation reduces to

$$\begin{aligned}(\frac{\partial^2\Pi}{\Pi})' - 2(\frac{\Pi'}{\Pi})(\frac{\partial^2\Pi}{\Pi}) &= \frac{\alpha^2}{2}(\tilde{\omega} + \frac{D}{x^3})\tilde{f} \\ \tilde{\omega} &= -2(\frac{\Pi'}{\Pi})^2(\frac{\Pi'}{\Pi} + \frac{3}{x}) \\ \tilde{f} &= \frac{\Pi'}{\Pi}(\frac{\Pi'}{\Pi} + \frac{2}{x})\end{aligned} \tag{12}$$

To preserve the Pontryagin index and finiteness of energy the boundary conditions on Π are taken to be the same as before i.e. $\Pi \longrightarrow \frac{\rho^2}{x^2}$ as $x \longrightarrow 0$ and $\Pi \longrightarrow 1 + \frac{\rho^2}{x^2}$ as $x \longrightarrow \infty$

At first sight the extreme non linearity of these equations seems intractable. One expects,however, that if one can solve it in the asymptotic regions $x \longrightarrow 0, x \longrightarrow \infty$ to obtain a solution with the same winding number as for $N_f = 0$ then a solution which interpolates between these regions may be obtained numerically. Evaluating the r.h.s of eqn(12) in the limit $x \longrightarrow 0$ one finds that the leading term is $-2\alpha^2(D-8)/(x^3\rho^2)$ while the l.h.s is less singular. Hence the choice $D = 8$ is confirmed. With $D = 8$ one finds that one can solve for the unknown coefficients in the deformed superpotential

$$\Pi_0 = 1 + \frac{\rho^2}{x^2} + a_1 x^2 + a_2 x^4 + \ldots \tag{13}$$

to get

$$a_1 = -\frac{\alpha^2}{\rho^4} \qquad a_2 = +\frac{2\alpha^2}{3\rho^6}(1 + \frac{9\alpha^2}{4\rho^2}) \tag{14}$$

and so on. Note that inspite of the nonlinearity of the fermion back reaction represented by the Chern-Simons term, the deformations are entirely non singular. Similarly in the $x \to \infty$ region

$$\Pi_\infty = 1 + \frac{\rho^2}{x^2} + \frac{b_1}{x^4} + \frac{b_2}{x^6} + \ldots \tag{15}$$

$$b_1 = \frac{\alpha^2\rho^2}{3} \qquad\qquad b_2 = \frac{\alpha^4\rho^2}{18} \tag{16}$$

A numerical integration of the equation in the intermediate region will be given elsewhere. The asymptotic behaviour of the electric field is

$$\begin{aligned} E^8_\mu &= -\partial_\mu A^8_0 = -\frac{\alpha}{2}(\tilde{\omega}_\mu + \frac{D\hat{x}_\mu}{x^3}) \\ &= -\frac{4\alpha}{x^3}\hat{x}_\mu + \frac{12\alpha\rho^4}{x^7}\hat{x}_\mu + O(x^{-9}) \end{aligned} \tag{17}$$

The electric charge is

$$Q^8 = \frac{1}{g_5^2}\int_{S_\infty} d\Sigma_\mu E^8_\mu = -\frac{N_f}{4\sqrt{3}} \tag{18}$$

The electric field near the origin is

$$E^8_\mu = -\frac{12\alpha}{\rho^4}x_\mu + O(x^3) \tag{19}$$

The magnetic field is

$$\begin{aligned} F^a_{\mu\nu} &= -\frac{4}{\rho^2}(\bar{\eta}_{a\mu\nu} + 2\hat{x}_\sigma\bar{\eta}_{a\sigma[\mu}\hat{x}_{\nu]}) + O(x^2) \qquad\qquad x \longrightarrow 0 \\ &= -\frac{4\rho^2}{x^4}(\bar{\eta}_{a\mu\nu} + 2\hat{x}_\sigma\bar{\eta}_{a\sigma[\mu}\hat{x}_{\nu]}) + O(x^{-6}) \qquad\qquad x \longrightarrow \infty \end{aligned} \tag{20}$$

The magnetic field is thus asymptotically *dipolar*.

The CS term does not contribute to the expression for the field energy :

$$\begin{aligned} \mathcal{E} &= \frac{8\pi\nu}{g_5^2} + \frac{3}{2g_5^2}\int d^4x(\frac{\partial^2\Pi}{\Pi})^2 - \frac{1}{2g_5^2}\int d^4x d^4y \rho_8(x)G_0(x,y)\rho_8(y) \\ \rho_8(x) &= -\frac{\alpha}{4}F^a_{\mu\nu}\tilde{F}^a_{\mu\nu} \qquad\qquad G_0(x,y) = -\frac{1}{4\pi^2(x-y)^2} \end{aligned} \tag{21}$$

Thus it is the sum of the original uncharged soliton energy plus a positive definite contribution from the electrostatic energy and another due to the deviation from self duality. The winding number is unaffected by the weak deformations exhibited in equations (13)-(16).

$$\begin{aligned}\nu &= \frac{1}{32\pi^2}\int d^4x F^a_{\mu\nu}\tilde{F}^a_{\mu\nu} = \frac{1}{16\pi^2}\int d^4x \partial_\mu \tilde{\omega}_\mu \\ &= \frac{1}{8}x^3 \hat{x}_\mu \tilde{\omega}_\mu \mid_0^\infty = 1\end{aligned} \tag{22}$$

This can be checked by transforming to a nonsingular gauge using the usual[12] transformation $U = (x^4 + i\vec{\tau}\cdot\vec{x})/\mid x\mid$. The above solutions are parametrized by an arbitrary scale parameter ρ. However the energy $\mathcal{E}$ is not independent of ρ . Thus one expects that the present solution, if stable at all, will relax to that value of ρ which minimizes the energy. To estimate this value one may approximate Π by Π_0 and Π_∞ in the regions $x \in [0,\rho], x \in [\rho,\infty]$ respectively to obtain

$$\mathcal{E} = \frac{8\pi^2}{g_5^2} + \frac{80\pi^2 g_5^6 N_f^4}{27(32\pi^2)^4\rho^4} + \frac{c g_5^2 N_f^2}{\rho^2} \tag{23}$$

Where the last term is the obvious dimensional estimate for the electrostatic energy in 4 space dimensions and c is a numerical constant. In the approximation we have used , it appears that our solution is unstable against growth of the free parameter ρ which will tend to ∞ so as to saturate the the self-duality lower bound on the energy. Note however that the corrections to the neutral soliton energy are $O(g_5^2)$ and $O(g_5^6)$ thus it quite possible that quantum corrections may stabilize ρ.

Since the electric and magnetic fields are orthogonal in the internal space, the Poynting vector, and therefore also the usual (Belinfante-Bessel-Hagen) expression for the the angular momentum of the field configuration, is zero.

It is easy to see that solutions with winding number K can be approximated by superpositions of Π_0 's centered at different x_i near those x_i and by $1 + \sum_i \rho_i^2/x^2 + O(x^{-4})$ in the $x \to \infty$ region. To generalize from $SU(3)$ to the $SU(N)$ case one simply replaces $\lambda^8 = \frac{1}{\sqrt{3}} diag(1,1,-2)$ by
$\{\lambda^{k^2-1} = \sqrt{\frac{2}{k(k-1)}} diag(1_{k-1}, 1-k, 0_{N-k}); k = 3, ..., N\}$ etc[13].

In conclusion we have demonstrated the existence of charged topological solitons in 5 dimensional YMCS theory. This extends results on charged vortices in 3 demensional YMCS theory and supports the conjecture that similar "charging" by the CS term occurs as well in higher odd dimensions. The generalizations of our solution to $M^4 \times S^1$ will be relevant to fleshing out at least the simplest example of the "syncyclonic scenario"[1]. The semiclassical quantization of of these objects is therefore studied in Ref.[9].

Acknowledgements: I am grateful to A.Khare, V.Soni, S.Wadia and R.Shankar for very useful discussions. I thank S.Kar for drawing my attention to M.Kobayashi's paper[2].

Note Added : We[13] have now shown that the different embeddings of $SU(2) \times U(1)^{N-2}$ in $SU(N)$ correspond to a ${}^N C_2$-plet of topological solitons. Moreover the "dibaryon embedding" i.e $SO(3) \times U(1)^{N-3}$ in $SU(N)$ gives *neutral* solutions for $N = 3$ and a ${}^N C_3$-plet for $N \geq 4$ (after modding out the action of certain $SU(3)$s).

Email:aulakh%iopb@shakti.ernet.in March 1992

References

1) C.S.Aulakh, *Syncyclons or Solitonic Signals from Extra Dimensions*, IP/BBSR/92-14, to be published in Mod. Phys. Lett **A**. See also Ref.[2].
2) A.Strominger Nucl. Phys. B 343 (1990) 167.
   M.Kobayashi, Prog. Theor. Phys. 74(1985)1139.
3) B.J.Harrington and H.K.Shepard, Phys. Rev. D17 (1978), 2122.
4) D.Gross, R.Pisarski, L.Yaffe, Rev. Mod. Phys. 53 (1981) 43.
5) A.Niemi and G.W.Semenoff, Physics Reports 135 (1986) 99 and references therein ;
   Phys. Rev. Lett. 51 (1983) 2077.
6) E.Witten,Phys.Lett. B86(1979)283.
7) E.Witten, Nucl. Phys. B223 (1983) 422.
   S.Jain and S.R.Wadia , Nucl. Phys. B258(1985) 713.
8) S.K.Paul and A.Khare Phys. Lett. B174(1986)420,182(1986),414.
   C.N.Kumar and A.Khare ,Phys.Lett. B 178(1986)385.
   H.J.De Vega and F.A.Schaposhnik ,Phys. Rev. Lett. 56(1986)2564.
9) C.S.Aulakh and V.Soni, *Collective Quantization of Topological Solitons in 4+1 Dimensional YMCS Theory*, IP/BBSR/92-37, to appear.
10) C.S.Aulakh, to appear.
11) G.'t Hooft (unpublished);Phys. Rev. D14(1976)3432.
    E.Corrigan and D.Fairlie, Phys. Lett. B67 (1977), 69.
    R.Jackiw, C.Nohl. and C.Rebbi, Phys. Rev. D15, (1977)1642.
12) R.Rajaraman, *Solitons and Instantons*, North Holland, 1982 and references therein.
13) C.S.Aulakh and V.Soni, *Topological Soliton Multiplets in 4+1 Dimensional YMCS Theory*, IP/BBSR/92-36, to appear.

Email:aulakh%iopb@shakti.ernet.in March 1992